

# **Tidal Stream Assessment Report for Bullfrog Creek 03100206000028 in Hillsborough County, Florida**

---

Date Assessed: June 5, 2012

Assessed by: David Eilers, Sawyer Masonjones, Daniel Dickinson

Reviewed by: Jim Griffin

## **INTRODUCTION**

This assessment was conducted to update existing physical and ecological data for Bullfrog Creek on the [Hillsborough County & City of Tampa Water Atlas](#). The Lake and Stream Assessment project is a collaborative effort between the University of South Florida's Center for Community Design and Research and Hillsborough County Stormwater Management Section. The project is funded by Hillsborough County. The project has, as its primary goal, the rapid assessing of up to 150 lakes and stream segments in Hillsborough County during a five-year period. The product of these investigations will provide the County, property owners and the general public a better understanding of the general health of Hillsborough County lakes and streams, in terms of shoreline development, water quality, morphology (bottom contour, volume, area, etc.) and the plant biomass and species diversity. These data are intended to assist the County and its citizens to better manage lakes and streams.



**Figure 1. Bullfrog Creek above US Highway 41.**

## BACKGROUND

This report focuses on Bullfrog Creek from Interstate 75 to the mouth in Tampa Bay. This region covers the tidal region of Bullfrog Creek, Reach 03100206000028, which consists of two stream segments<sup>i</sup> (please see maps, Figures 2-11). The lower segment near the mouth at Tampa Bay is characterized by a meandering channel through a mangrove dominated estuary. The upper segment is characterized by a narrower channel winding through more typical freshwater stream vegetation.

**The first section** of the report provides the results of the overall morphological assessment of the stream. Primary data products include: a contour (bathymetric) map of the stream, area, volume and depth statistics, and the water level at the time of assessment. These data are useful for evaluating trends and for developing management actions such as plant management where depth and stream volume are needed.

**The second section** provides the results of the vegetation assessment conducted on the stream. These results can be used to better understand and manage vegetation in the stream. A list is provided with the different plant species found at various sites along the stream. Potentially invasive, exotic (non-native) species are identified in a plant list and the percent of exotics is presented in a summary table. Watershed values provide a means of reference.

**The third section** provides the results of the water quality sampling of the stream. Both field data and laboratory data are presented. The stream water quality is assessed based on the Water Quality Standards for Streams that were fully approved on March 15, 2013. Please see a discussion of these standards and the approach used for tidal streams in the Stream Assessment Notes at the end of this report.

The intent of this assessment is to provide a starting point from which to track changes in the stream, and where previous comprehensive assessment data is available, to track changes in the stream's general health. These data can provide the information needed to determine changes and to monitor trends in physical condition and ecological health of the stream.

### Section 1: Stream Morphology

**Bathymetric Map**<sup>ii</sup>. Table 1 provides the stream's morphologic parameters in various units. The bottom of the stream was mapped using a Lowrance LCX 28C HD or HDS 5 with Wide Area Augmentation System (WAAS)<sup>iii</sup> enabled Global Positioning System (GPS) with fathometer (bottom sounder) to determine the boat's position, and bottom depth in a single measurement. The result is an estimate of the stream's area, mean and maximum depths, and volume and the creation of a bottom contour map (Figure 2). Besides pointing out the deeper fishing holes in the stream, the morphologic data derived from this part of the assessment can be valuable to overall management of the stream vegetation as well as providing flood storage data for flood models.

---

<sup>i</sup> A stream segment is a stream length that can be conveniently mapped and assessed during a single assessment period.

<sup>ii</sup> A bathymetric map is a map that accurately depicts all of the various depths of a water body. An accurate bathymetric map is important for effective herbicide application and can be an important tool when deciding which form of management is most appropriate for a water body. Stream volumes, hydraulic retention time and carrying capacity are important parts of stream management that require the use of a bathymetric map.

<sup>iii</sup> WAAS is a form of differential GPS (DGPS) where data from 25 ground reference stations located in the United States receive GPS signals from GPS satellites in view and retransmit these data to a master control site and then to geostationary satellites. For more information, see end note 3.

**Table 1. Stream Morphologic Data (Area, Depth and Volume)**

Parameter	Feet	Meters	Acres	Acre-Ft	Gallons
Surface Area (sq)	4,787,523.8	444,775.5	109.9		
Mean Depth	3.22	0.98			
Maximum Depth	11.44	3.49			
Volume (cubic)	39,993,618	1,132,493.15		918.13	299,175,111.5
Gauge (USGS 02300700)	18.61	5.67			





Figure 2. 2012 Bullfrog Creek 2-Foot Bathymetric Contour Map Overview





**Figure 3 Bullfrog Creek 2012 Bathymetric Contour Inset Map 1**





Figure 4 Bullfrog Creek 2012 Bathymetric Contour Inset Map 2





Figure 5 Bullfrog Creek 2012 Bathymetric Contour Inset Map 3





**Figure 6 Bullfrog Creek 2012 Bathymetric Contour Inset Map 4**





**Figure 7 Bullfrog Creek 2012 Bathymetric Contour Inset Map 5**






Figure 8 Bullfrog Creek 2012 Bathymetric Contour Inset Map 6






Figure 9 Bullfrog Creek 2012 Bathymetric Contour Inset Map 7

## Section 2: Stream Ecology (Vegetation)

The stream's apparent vegetative cover and shoreline detail are evaluated using the latest stream aerial photograph as shown in and by use of WAAS-enabled GPS. Submerged vegetation is determined from the analysis of bottom returns from the Lowrance 28c HD or HDS 5 combined GPS/fathometer described earlier. As depicted in Figure 10-11, 32 vegetation regions have been assessed for in ~250 meter regions measured from the center of the stream. The vegetation assessment regions are set up from the upstream extent and work to the downstream extent. The region beginning and ending points are set using GPS and then loaded into a GIS mapping program (ArcGIS) for display. Each region is sampled in the three primary vegetative zones (emergent, submerged and floating)<sup>iv</sup>. The latest high-resolution aerial photos are used to provide shore details (docks, structures, vegetation zones) and to calculate the extent of surface vegetation coverage. The primary indices of submerged vegetation cover and biomass for the stream, percent area coverage (PAC) and percent volume inhabited (PVI), are determined by transiting the stream by boat and employing a fathometer to collect "hard and soft return" data. These data are later analyzed for presence and absence of vegetation and to determine the height of vegetation if present. The PAC is determined from the presence and absence analysis of 100 sites in the stream and the PVI is determined by measuring the difference between hard returns (stream bottom) and soft returns (top of vegetation) for sites (within the 100 analyzed sites) where plants are determined present.

The data collected during the site vegetation sampling include vegetation type, exotic vegetation, predominant plant species and submerged vegetation biomass. The total number of species from all sites is used to approximate the total diversity of aquatic plants and the percent of invasive-exotic plants on the stream (

Table 2). The Watershed value in Table 2 only includes lakes and streams sampled during the stream assessment project begun in May of 2006. These data will change as additional lakes and streams are sampled.

---

<sup>iv</sup> See end note 3.






Figure 11 Bullfrog Creek Vegetation Assessment Map Regions 13-32

Table 3 through Table 5 detail the results from the 2012 aquatic plant assessment for the stream. These data are determined from the 32 regions used for intensive vegetation surveys. The tables are divided into Floating Leaf, Emergent and Submerged plants and contain the plant code, species, common name and the calculated percent occurrence (number sites species is found/number of sites) and type of plant (Native, Non-Native, Invasive, Pest). In the “Type” category, the codes N and E0 denote species native to Florida. The code E1 denotes Category I invasive species, as defined by the [Florida Exotic Pest Plant Council](#) (FLEPPC); these are species “that are altering native plant communities by displacing native species, changing community structures or ecological functions, or hybridizing with natives.” The code E2 denotes Category II invasive species, as defined by FLEPPC; these species “have increased in abundance or frequency but have not yet altered Florida plant communities to the extent shown by Category I species.” Use of the term invasive indicates the plant is commonly considered invasive in this region of Florida. The term “pest” indicates a plant (native or non-native) that has a greater than 55% occurrence in the stream and is also considered a problem plant for this region of Florida, or is a non-native invasive that is or has the potential to be a problem plant in the stream and has at least 40% occurrence. These two terms are somewhat subjective; however, they are provided to give stream property owners some guidance in the management of plants on their property.

Please remember that to remove or control plants in a wetland (stream shoreline) in Hillsborough County the property owner must secure an [Application To Perform Miscellaneous Activities In Wetlands](#) permit from the [Environmental Protection Commission of Hillsborough County](#) and for management of in-stream vegetation outside the wetland fringe (for streams with an area greater than ten acres), the property owner must secure a [Florida Department of Environmental Protection Aquatic Plant Removal Permit](#).

**Table 2. Total Diversity, Percent Exotics, and Number of Pest Plant Species**

Parameter	Lake	Watershed
Number of Vegetation Assessment Sites	32	32
Total Plant Diversity (# of Taxa)	98	98
% Non-Native Plants	30	26
Total Pest Plant Species	1	1





**Figure 10. Bullfrog Creek Vegetation Assessment Map regions 1-13**






Figure 11 Bullfrog Creek Vegetation Assessment Map Regions 13-32





**Figure 12.** West Indian Manatee's were observed during the assessment in the upper regions of Bullfrog Creek

**Table 3. List of Emergent Zone Aquatic Plants Found**

Plant Species Code	Scientific Name	Common Name	Percent Occurrence	Type
STS	<i>Schinus terebinthifolius</i>	Brazilian Pepper	78%	E1, P
SMR	<i>Sabal minor</i>	Dwarf Palmetto; Bluestem Palm	65%	N, E0
QVA	<i>Quercus virginiana</i>	Virginia Live Oak	65%	N, E0
LAG	<i>Laguncularia racemosa</i>	White Mangrove	62%	N, E0
RZM	<i>Rhizophora mangle</i>	Red Mangrove	62%	N, E0
ADM	<i>Acrostichum danaeifolium</i>	Giant Leather Fern	56%	N, E0
AVG	<i>Avicennia germinans</i>	Black Mangrove	50%	N, E0
LEL	<i>Leucaena leucocephala</i>	White Leadtree	50%	E2,P
VRA	<i>Vitis rotundifolia</i>	Muscadine Grape	50%	N, E0
JRO	<i>Juncus roemerianus</i>	Needle Rush, Black Rush	40%	N, E0
TYP	<i>Typha spp.</i>	Cattails	34%	N, E0
XAA	<i>Ximenia americana</i>	Tallow Wood, Hog Plum	34%	N, E0
SPO	<i>Sabal palmetto</i>	Sabal Palm, Cabbage Palm	31%	N, E0
QLA	<i>Quercus laurifolia</i>	Laurel Oak; Diamond Oak	31%	N, E0
DBA	<i>Dioscorea bulbifera</i>	Air Potato	31%	E1
CAM	<i>Crinum americanum</i>	Swamp lily	28%	N, E0
PAN	<i>Panicum spp.</i>	Panic Grasses	28%	E0
WAX	<i>Myrica cerifera</i>	Southern Bayberry; Wax Myrtle	28%	N, E0
WTA	<i>Sphagneticola trilobata</i>	Creeping Oxeye; Wedelia	28%	E2
CEA	<i>Colocasia esculenta</i>	Wild Taro	25%	E1



Plant Species Code	Scientific Name	Common Name	Percent Occurrence	Type
CLA	<i>Casuarina equisetifolia</i>	Australian Pine	25%	E1
APS	<i>Alternanthera philoxeroides</i>	Alligator Weed	21%	E2
BHA	<i>Baccharis halimifolia</i>	Groundsel Tree; Sea Myrtle	21%	N, E0
RBA	<i>Ruellia simplex</i>	Britton's Wild Petunia	21%	E1
PRA	<i>Pluchea baccharis</i>	Rosy Camphorweed	21%	N, E0
PRS	<i>Panicum repens</i>	Torpedo Grass	21%	E1
SAM	<i>Sambucus nigra subsp. Canadensis</i>	Elderberry	21%	N, E0
RF	<i>Osmunda regalis var. spectabilis</i>	Royal Fern	18%	N, E0
PQA	<i>Parthenocissus quinquefolia</i>	Virginia Creeper, Woodbine	18%	N, E0
SMI	<i>Smilax spp.</i>	Catbriar, Greenbriar	18%	N, E0
PIN	<i>Pinus spp.</i>	Pine Tree	18%	N, E0
OCA	<i>Osmunda cinnamomea</i>	Cinnamon Fern	18%	N, E0
LPA	<i>Ludwigia peruviana</i>	Peruvian Primrosewillow	18%	E1
MAH	<i>Melia azedarach</i>	Chinaberry tree	18%	E2
EUP	<i>Eupatorium capillifolium</i>	Dog Fennel	18%	N, E0
ACS	<i>Symphyotrichum carolinianum</i>	Climbing Aster	18%	N, E0
BRP	<i>Broussonetia papyrifera</i>	Paper Mulberry	18%	E2
BAA	<i>Bidens alba</i>	White Beggar-ticks, Romerillo	18%	N, E0
BAM	<i>Bambusa spp.</i>	Bamboo	18%	E0
CAL	<i>Callicarpa americana</i>	Beautyberry	18%	N, E0
CAQ	<i>Carya aquatica</i>	Water Hickory	18%	N, E0
ENT	<i>Enterolobium contortisiliquum</i>	Earpod Tree	15%	E2
ERH	<i>Erechtites hieraciifolia</i>	Fireweed	15%	N,

Plant Species Code	Scientific Name	Common Name	Percent Occurrence	Type
				E0
BMA	<i>Urochloa mutica</i>	Para Grass	15%	E1
LJM	<i>Lygodium japonicum</i>	Japanese Climbing Fern	15%	E1
NEA	<i>Nephrolepis exaltata</i>	Sword Fern, Wild Boston Fern	15%	N, E0
QNA	<i>Quercus nigra</i>	Water Oak	15%	N, E0
RVS	<i>Rumex verticillatus</i>	Swamp Dock	15%	N, E0
SPM	<i>Syngonium podophyllum</i>	Nepthytis, Arrowhead Vine, American Evergreen	12%	E1
OXS	<i>Tiedemannia filiformis</i>	Water cowbane	12%	N, E0
CIS	<i>Cyperus involucratus</i>	Umbrella Flat Sedge	12%	E2
HYE	<i>Hydrocotyle umbellata</i>	Manyflower Marshpennywort, Water Pennywort	12%	N, E0
IFS	<i>Iva frutescens</i>	Bigleaf Sumpweed, Marsh Elder	12%	N, E0
BOC	<i>Boehmeria cylindrica</i>	Bog Hemp, False Nettle	12%	N, E0
ADX	<i>Arundo donax</i>	Giant Reed	12%	E0
DEM	<i>Dalbergia ecastaphyllum</i>	Coinvine	12%	N, E0
DIO	<i>Diospyros virginiana</i>	Common Persimmon	9%	N, E0
COM	<i>Commelina spp.</i>	Dayflower	9%	N, E0
AAA	<i>Ampelopsis arborea</i>	Peppervine	9%	N, E0
JVA	<i>Juniperus virginiana</i>	Red cedar	9%	N, E0
MSS	<i>Mikania scandens</i>	Climbing Hempvine	9%	N, E0
JES	<i>Juncus effusus subsp. Solutus</i>	Soft Rush	9%	N, E0
PPP	<i>Pleopeltis polypodioides var.</i>	Resurrection Fern	9%	N,



Plant Species Code	Scientific Name	Common Name	Percent Occurrence	Type
	<i>michauxiana</i>			E0
SSS	<i>Solidago sempervirens</i>	Seaside Goldenrod	9%	N, E0
TRD	<i>Tripsacum dactyloides</i>	Eastern Gama Grass	6%	N, E0
TRS	<i>Toxicodendron radicans</i>	Poison Ivy	6%	N, E0
SHA	<i>Sesbania herbacea</i>	Danglepod Sesban	6%	N, E0
ULA	<i>Urena lobata</i>	Caesar's-weed	6%	E1
ITE	<i>Itea virginica</i>	Virginia Willow; Virginia Sweetspire	6%	N, E0
HGS	<i>Hibiscus grandiflorus</i>	Swamp Rosemallow, Swamp Hibiscus	6%	N, E0
ACE	<i>Acer rubrum</i>	Southern Red Maple	6%	N, E0
BCS	<i>Baccharis spp.</i>	Falsewillow	6%	N, E0
BVE	<i>Blutaparion vermiculare</i>	Silverhead, Saltweed	6%	N, E0
ANT	<i>Antigonon leptopus</i>	Coralvine	6%	E2
BLS	<i>Blechnum serrulatum</i>	Swamp fern, Toothed Midsorus Fern	6%	N
COS	<i>Cephalanthus occidentalis</i>	Buttonbush	6%	N, E0
CDS	<i>Cyperus difformis</i>	Variable Flatsedge	6%	E0
COC	<i>Coccoloba uvifera</i>	Seagrape	3%	N, E0
CFO	<i>Cornus foemina</i>	Swamp Dogwood, Stiff Dogwood	3%	N, E0
CRS	<i>Campsis radicans</i>	Trumpet Vine	3%	N, E0
DSO	<i>Dalbergia sissoo</i>	Indian Rosewood	3%	E2
AJN	<i>Albizia julibrissin</i>	Silk Tree, Mimosa	3%	E1
EWI	<i>Echinochloa walteri</i>	Coast Cockspur Grass (hairy)	3%	N, E0
GTM	<i>Galium tinctorium</i>	Marsh Bedstraw	3%	N,

Plant Species Code	Scientific Name	Common Name	Percent Occurrence	Type
				E0
ICE	<i>Ilex cassine</i>	Dahoon Holly	3%	N, E0
MUS	<i>Musa spp.</i>	Banana Tree	3%	E0
RCS	<i>Ricinus communis</i>	Castorbean	3%	E2
PBA	<i>Persea borbonia</i>	Redbay	3%	N, E0
PEP	<i>Persea palustris</i>	Swampbay	3%	N, E0
PHS	<i>Polygonum hydropiperoides</i>	Mild Waterpepper; Swamp Smartweed	3%	N, E0
SLA	<i>Sagittaria lancifolia</i>	Duck Potato	3%	N, E0
SCA	<i>Salix caroliniana</i>	Carolina Willow	3%	N, E0
SAC	<i>Sacciolepis striata</i>	American Cupscale	3%	N, E0
TDM	<i>Taxodium distichum</i>	Bald Cypress	3%	N, E0





**Figure 13. Typical emergent vegetation community on Bullfrog Creek in the estuarine region**

**Table 4. List of Submerged Zone Aquatic Plants Found.**

Plant Species Code	Scientific Name	Common Name	Percent Occurrence	Type
BMI	<i>Bacopa monnieri</i>	Common Bacopa	12%	N, E0
SFE	<i>Syringodium filiforme</i>	Manateeegrass	9%	N, E0
ULV	<i>Ulva spp.</i>	Sea Lettuce	3%	N, E0
RUP	<i>Ruppia maritima</i>	Widgeongrass	3%	N, E0





**Figure 14. Typical emergent vegetation community on Bullfrog Creek in the freshwater portion**

**Table5. List of All Plants and Sample Sites**

Plant Common Name	Found at Sample Sites	Percent Occurrence	Growth Type
Brazilian Pepper	1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23,24,29	78	Emergent
Dwarf Palmetto; Bluestem Palm	1,3,4,5,6,7,8,9,11,12,13,14,15,16,17,18,19,20,21,23,29	65	Terrestrial
Virginia Live Oak	2,3,4,5,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,24	65	Terrestrial
Red Mangrove	12,13,14,15,16,17,18,20,21,22,23,24,25,26,27,28,29,30,31,32	62	Terrestrial
White Mangrove	12,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32	62	Terrestrial
Giant Leather Fern	1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,21,30	56	Emergent
Black Mangrove	14,15,17,18,21,22,23,24,25,26,27,28,29,30,31,32	50	Terrestrial
Muscadine Grape	1,2,3,4,5,6,7,8,9,10,11,12,14,15,16,20	50	Emergent
White Leadtree	2,3,4,5,6,7,8,9,11,12,13,14,16,21,22,24	50	Terrestrial
Needle Rush, Black Rush	11,12,14,15,16,17,18,19,20,21,26,29,30	40	Emergent
Cattails	1,5,7,9,10,11,12,13,14,15,16	34	Emergent
Tallow Wood, Hog Plum	1,3,7,8,9,10,11,12,15,16,17	34	Terrestrial
Air Potato	1,2,3,4,5,6,7,8,10,13	31	Emergent
Laurel Oak; Diamond Oak	1,2,3,4,7,8,9,12,16,17	31	Emergent
Sabal Palm, Cabbage Palm	1,4,5,7,9,10,11,16,17,30	31	Terrestrial
Creeping Oxeye; Wedelia	1,3,4,5,7,8,10,11,16	28	Emergent
Panic Grasses	1,2,3,4,5,6,7,8,12	28	Emergent
Southern Bayberry; Wax Myrtle	1,3,4,5,8,9,10,11,12	28	Emergent
Swamp lily	1,7,8,9,10,11,12,15,16	28	Emergent
Australian Pine	7,9,10,13,15,22,23,24	25	Emergent
Wild Taro	1,2,3,4,5,6,7,10	25	Emergent
Alligator Weed	5,6,7,8,9,10,11	21	Emergent
Britton's Wild Petunia	1,2,3,4,5,8,9	21	Terrestrial
Elderberry	1,2,3,4,5,6,7	21	Emergent
Groundsel Tree; Sea Myrtle	6,7,10,11,13,14,22	21	Emergent
Rosy Camphorweed	2,3,4,5,6,7,11	21	Emergent
Torpedo Grass	9,10,11,12,13,16,22	21	Emergent
Bamboo	1,3,4,5,12,19	18	Emergent
Beautyberry	2,4,5,7,10,11	18	Emergent
Catbriar, Greenbriar	1,5,7,8,11,16	18	Emergent
Chinaberry tree	2,3,5,6,8,12	18	Emergent
Cinnamon Fern	1,2,3,4,5,6	18	Emergent



Plant Common Name	Found at Sample Sites	Percent Occurrence	Growth Type
Climbing Aster	2,3,4,5,6,9	18	Emergent
Dog Fennel	1,2,3,4,7,13	18	Emergent
Paper Mulberry	2,3,7,8,9,13	18	Emergent
Peruvian Primrosewillow	1,2,3,4,5,8	18	Emergent
Pine Tree	11,12,20,21,23,29	18	Emergent
Royal Fern	1,3,4,5,6,7	18	Emergent
Virginia Creeper, Woodbine	1,2,5,6,8,13	18	Emergent
Water Hickory	1,2,3,4,9,11	18	Emergent
White Beggar-ticks, Romerillo	1,2,3,5,7,8	18	Terrestrial
Earpod Tree	3,4,5,8,16	15	Emergent
Fireweed	1,2,4,6,7	15	Terrestrial
Japanese Climbing Fern	1,3,4,6,7	15	Terrestrial
Para Grass	1,2,3,4,8	15	Emergent
Swamp Dock	1,4,5,6,9	15	Emergent
Sword Fern, Wild Boston Fern	3,5,6,8,10	15	Terrestrial
Water Oak	5,6,7,8,10	15	Emergent
Bigleaf Sumpweed, Marsh Elder	15,16,17,23	12	Terrestrial
Bog Hemp, False Nettle	1,2,3,5	12	Emergent
Coinvine	8,12,17,22	12	Terrestrial
Common Bacopa	4,6,8,10	12	Submersed
Giant Reed	2,3,5,12	12	Terrestrial
Manyflower Marshpennywort, Water Pennywort	1,2,3,4	12	Emergent
Nephthytis, Arrowhead Vine, American Evergreen	2,3,4,10	12	Terrestrial
Umbrella Flat Sedge	3,7,9,12	12	Emergent
Water cowbane	2,4,5,7	12	Terrestrial
Climbing Hempvine	1,2,5	9	Emergent
Common Persimmon	1,6,10	9	Emergent
Dayflower	1,2,4	9	Emergent
Manateegrass	30,31,32	9	Submersed
Peppervine	1,2,3	9	Emergent
Red cedar	14,20,21	9	Terrestrial

Plant Common Name	Found at Sample Sites	Percent Occurrence	Growth Type
Resurrection Fern	1,2,4	9	Emergent
Seaside Goldenrod	4,11,16	9	Terrestrial
Soft Rush	6,7,8	9	Emergent
Buttonbush	1,10	6	Emergent
Caesar's-weed	1,2	6	Emergent
Coralvine	1,8	6	Terrestrial
Danglepod Sesban	1,2	6	Emergent
Eastern Gama Grass	4,5	6	Emergent
Falsewillow	16,17	6	Emergent
Poison Ivy	1,5	6	Emergent
Silverhead, Saltweed	12,16	6	Terrestrial
Southern Red Maple	3,7	6	Emergent
Swamp fern, Toothed Midsorus Fern	7,8	6	Emergent
Swamp Rosemallow, Swamp Hibiscus	3,5	6	Emergent
Variable Flatsedge	5,6	6	Emergent
Virginia Willow; Virginia Sweetspire	6,7	6	Emergent
American Cupscale	1	3	Emergent
Bald Cypress	3	3	Emergent
Banana Tree	3	3	Emergent
Carolina Willow	8	3	Emergent
Castorbean	1	3	Terrestrial
Coast Cockspur Grass (hairy)	4	3	Emergent
Dahoon Holly	10	3	Emergent
Duck Potato	4	3	Emergent
Indian Rosewood	3	3	Terrestrial
Marsh Bedstraw	1	3	Emergent
Mild Waterpepper; Swamp Smartweed	1	3	Emergent
Redbay	10	3	Emergent
Sea Lettuce	24	3	Submersed
Seagrape	23	3	Terrestrial
Silk Tree, Mimosa	5	3	Emergent
Swamp Dogwood, Stiff Dogwood	5	3	Emergent
Swampbay	6	3	Emergent



Plant Common Name	Found at Sample Sites	Percent Occurrence	Growth Type
<b>Trumpet Vine</b>	<b>4</b>	<b>3</b>	<b>Emergent</b>
<b>Widgeongrass</b>	<b>28</b>	<b>3</b>	<b>Submersed</b>

## Discussion of Vegetation Assessment Results

The highest diversity of species found in the Bullfrog Creek study area was in the natural region of Bullfrog Creek region 1. In this region 45 species were identified. 35.5% of these species were non-native to Florida and 1 species was considered pest plants for this region. This region is located furthest upstream of all regions in the assessment. In this region salinity and tidal influence are minimal allowing more species to colonize the shorelines including many species typical of freshwater riparian zones and non-native invasive species. Regions 25, 27 and 28 had the lowest number of species identified with 3 species but also the lowest number of non-native species with no non-native species and no pest plant species present. These regions are located near the mouth of Bullfrog Creek and receive daily variations of salinity and tidal stresses. This area is dominated by salt tolerant plants such as the three species of mangroves. At the mouth at Tampa Bay, Bullfrog Creek is a predominately a mangrove lined estuary.

## Section 3: Long-term Ambient Water Chemistry

A critical element in any stream assessment is the long-term water chemistry data set. These data are obtained from several data sources that are available to the Water Atlas and are managed in the Water Atlas Data Download and graphically presented on the water quality page for streams in Hillsborough County. The Bullfrog Creek Water Quality Page can be viewed at <http://www.hillsborough.wateratlas.usf.edu/river/waterquality.asp?wbodyid=18&wbodyatlas=river>.

A primary source of stream water chemistry in Hillsborough County is the Routine Monitoring Sampling by the Hillsborough County Environmental Protection Commission. Other source data are used as available; however these data can only indicate conditions at time of sampling.

These data are displayed and analyzed on the Water Atlas as shown in Figures 15, 16, 17 and 18 for Bullfrog Creek. The figures are graphs of: (1) the overall water quality index (WQI)<sup>v</sup>, which is a method commonly used to characterize the productivity of a stream, and may be thought of as a stream's ability to support plant growth and a healthy food source for aquatic life; (2) the chlorophyll a concentration, which indicates the stream's algal concentration; (3) the stream's Secchi Disk depth which is a measure of water visibility and depth of light penetration and (4) the salinity which can be used to determine the extent of the tidal reach and measured as conductivity in micro Siemens per centimeter ( $\mu\text{S}/\text{cm}$ ) and commonly expressed as salinity in parts per million (ppt) and chloride concentration in milligrams/liter ( $\text{mg}/\text{L}$ )<sup>vi</sup>. These data are used to evaluate a stream's ecological health and to provide a method of ranking streams. These and other parameters (primary nutrients) are used by the US Environmental Protection Agency (USEPA) and the Florida Department of Environmental Protection (FDEP) to determine the level of impairment of a freshwater stream and tidal stream. The chlorophyll a and Secchi Disk depth graphs include benchmarks which indicate the median values for the various parameters for a large number of streams in Florida expressed as percentiles.

---

<sup>v</sup> See WQI discussion in Stream Assessment Notes at end of report.

<sup>vi</sup> Please see discussion of salinity in Stream Assessment notes at end of report.






Figure 15. Recent Water Quality Index (WQI) graph for Bullfrog Creek<sup>vii</sup>

<sup>vii</sup> Graph source: Hillsborough County Water Atlas. For an explanation of the Good, Fair and Poor benchmarks, please see the notes at the end of this report. For the latest data go to: [http://www.hillsborough.wateratlas.usf.edu/graphs20/graph\\_it.aspx?wbodyid={WBODY\\_ID}&data=WQI&datatype=WQ&waterbodyatlas=river&ny=10&bench=1](http://www.hillsborough.wateratlas.usf.edu/graphs20/graph_it.aspx?wbodyid={WBODY_ID}&data=WQI&datatype=WQ&waterbodyatlas=river&ny=10&bench=1)




Figure 16. Recent Chlorophyll a graph for Bullfrog Creek<sup>viii</sup>

<sup>viii</sup> Graph Source: Hillsborough County Water Atlas. For the latest data go to [http://www.hillsborough.wateratlas.usf.edu/graphs20/graph\\_it.aspx?wbodyid={WBODY\\_ID}&data=Chla\\_ugl&datatype=WQ&waterbodyatlas=river&ny=10&bench=1](http://www.hillsborough.wateratlas.usf.edu/graphs20/graph_it.aspx?wbodyid={WBODY_ID}&data=Chla_ugl&datatype=WQ&waterbodyatlas=river&ny=10&bench=1)







Figure 17. Recent Secchi Disk graph for Bullfrog Creek<sup>ix</sup>

<sup>ix</sup> Graph Source: Hillsborough County Water Atlas. For the latest data go to [http://www.hillsborough.wateratlas.usf.edu/graphs20/graph\\_it.aspx?wbodyid={WBODY\\_ID}&data=secchi\\_ft&datatype=WQ&waterbodyatlas=stream&ny=10&bench=1](http://www.hillsborough.wateratlas.usf.edu/graphs20/graph_it.aspx?wbodyid={WBODY_ID}&data=secchi_ft&datatype=WQ&waterbodyatlas=stream&ny=10&bench=1)



**Figure 18. Graph of Salinity in Bullfrog Creek**

**Stream Numeric Nutrient Criteria.** November 30, 2012 the USEPA accepted the majority of the FDEP proposed NNCs which included an NNC for streams. In its proposed criteria, FDEP stated that tidal reaches of streams should be covered under the Florida Narrative Criteria<sup>x</sup>. However; tidal streams were not accepted at that time. On March 15, 2013, the USEPA also accepted the tidal creek criteria. The narrative criterion requires that the balance in natural populations of aquatic flora and fauna is maintained. For a tidal creek this can be interpreted maintaining the flora and fauna in the stream and the estuary reach to which the flows. A Tidal Creek Study will be conducted in the fall of 2013 with the goal of developing a proposed procedure for evaluating tidal creeks and for establishing numeric nutrient requirements. In the absence of an approved approach for assessing tidal creeks, the Lake and Stream Assessment program has adopted an methodology that was proposed by the USEPA in their technical support document for Florida numeric nutrient criteria which includes a section for tidal creeks published November 30, 2012 (please see excerpt and reference in Stream Assessment Notes at the end of this report). The methodology proposes two approaches which consider the upstream (freshwater segment and the downstream (estuarine segment). The methodology puts forward two approaches.

<sup>x</sup> Narrative Criteria states: 62-302-530(47)(b), Florida Administrative Code (F.A.C.), provides that “[i]n no case shall nutrient concentrations of a body of water be altered so as to cause an imbalance in natural populations of aquatic flora or fauna.”

- The first divides the segment by the mean chloride concentration of a segment<sup>xi</sup>. For segments that have a mean chloride concentration of greater than or equal to 1,500 mg/L, the estuarine criteria is used, and for those less than this value, the freshwater criteria is used.
- The second approach is a bit more complicated, it sets as the boundary conditions the approved numeric nutrient condition for the freshwater stream segment and the approved NNC for the estuarine reach and employs a relationship with salinity to calculate tidal creek NNC. The formula is then:

$$C_{TC} = C_{FW} + (S_{TC} - S_{FW}) \times (C_{Est} - C_{FW} / S_{Est} - S_{FW}) \text{ (Equation 1)}$$

where:

$C_{TC}$  = nutrient criterion for tidal creek segment

$C_{FW}$  = nutrient criterion for adjoining/upstream freshwater segment

$C_{Est}$  = nutrient criterion for adjoining estuarine segment

$S_{TC}$  = mean salinity for tidal creek segment

$S_{FW}$  = mean salinity for adjoining/upstream freshwater segment

$S_{Est}$  = mean salinity for adjoining estuarine segment

The NNC for freshwater streams is provided in the Stream Assessment Notes at the end of this report, and for the Tampa Bay area (considered West Central) total phosphorous must be less than or equal to 0.49 mg/L and total nitrogen must be less than or equal to 1.65 mg/L to meet the criteria and chlorophyll a must be at or below 20 µg/L not be considered impaired. The estuarine criteria for Tampa Bay are provided below.

Nutrient Water-shed Region	Total Phosphorus (tons/million m <sup>3</sup> )	Total Phosphorus (mg/L)	Total Nitrogen (tons/million m <sup>3</sup> )	Total Nitrogen (mg/L)	Chlorophyll a (µg/L)
Old Tampa Bay	0.23	0.21	1.08	0.98	9.3
Hillsborough Bay	1.28	1.16	1.62	1.47	15
Middle Tampa Bay	0.24	0.218	1.24	1.13	8.5
Lower Tampa Bay	0.14	0.127	0.97	0.89	5.1

Where the conversion 1 ton / (million cubic meters) = 0.907 mg/L is used to convert to commonly used values.

For Bullfrog Creek, a tidal creek that flows into Hillsborough Bay has three long-term data stations whose three-year geometric mean for Chlorophyll a, Total Nitrogen, Total Phosphorus and Chloride are as shown below. Note: 1 ton / (million cubic meter) = 0.907 mg/L

---

<sup>xi</sup> The 1,500 mg/L chloride threshold is used to define waters as *predominantly freshwater* or *predominantly marine water* [F.A.C. 62-302.200(22) and 62-302.200(23)].




**Table 3. Bullfrog Creek Data (2010-2012) As Geometric Mean of Values**

Stations/Parameter	132	167	144	NNC Fresh Water	NNC Salt Water (Hillsborough Bay)
Chlorophyll a (ug/L)	5.7	3.4	10.3	≤ 20	≤ 15
Nitrogen (mg/L)	1.664	0.837	0.824	≤ 1.65	≤ 1.46
Phosphorous (mg/L)	0.188	0.221	0.268	≤ 0.49	≤ 1.16
Chloride (mg/L)	158.99	165.86	6,614.52	≤1,500	>1,500


Note: Fresh Water is less than 1,500 mg/L Chloride

The geometric mean of Chloride data for all stations in Hillsborough Bay is 20,497.21 mg/L Chloride.


Using the first approach, Bullfrog Creek stations 132 and 167, which are both east of I-75 and are fresh water stations (Chloride less than or equal to 1,500 mg/L), would use the freshwater standards, and station 144, with chloride significantly above the freshwater standard, would use the estuarine standard. The problem with this approach is that we do not really know where the tidal portion begins. As shown in **Error! Reference source not found.**, both freshwater stations would meet Chlorophyll a NNC; but as seen in **Error! Reference source not found.**, Station 132 fails the nitrogen NNC. All three stations pass the Total Phosphorous criteria.



**Figure 19 Chlorophyll a sample values for three Bullfrog Creek stations 2010-2012**



**Figure 20 Total Nitrogen sample values for three Bullfrog Creek Stations 2010-2012**



**Figure 21. Total Phosphorous values for three Bullfrog Creek stations 2010-2012**

The second approach allows the estimation of the tidal section and estimation of the NNC for this section. Using the geometric mean for salinity for tidal portion of the creek of 25 ppt (station 144) and the mean for the freshwater segment of 0.22 (stations 132 and 167) and the NNCs for West-Central freshwater streams and for Hillsborough Bay and Equation 1 below, the calculated Total Phosphorus NNC for the tidal portion of Bullfrog Creek would be 0.88 mg/L and the calculated Total Nitrogen NNC would be 1.55 mg/L. With Figure 2 as a reference, the tidal portion of Bullfrog Creek would have two violations of the NNC based on the first approach and one based on the second approach. In either case the tidal portion would not be impaired for nitrogen. There would be no violations for TP under either approach.

$$C_{TC} = C_{FW} + (S_{TC} - S_{FW}) \times ((C_{Est} - C_{FW}) / (S_{Est} - S_{FW}))$$

$$C_{TCp} = 0.49 \text{ mg/L} + (14.8 - 0.22) \times ((1.16 \text{ mg/L} - 0.49 \text{ mg/L}) / (25 - 0.22)) = 0.88 \text{ mg/L Total Phosphorus}$$

$$C_{TCn} = 1.65 \text{ mg/L} + (14.8 - 0.22) \times ((1.47 \text{ mg/L} - 1.65 \text{ mg/L}) / (25 - 0.22)) = 1.55 \text{ mg/L Total Nitrogen}$$

$C_{TC}$  = nutrient criterion for tidal creek segment

$C_{FW}$  = nutrient criterion for adjoining/upstream freshwater segment

$C_{Est}$  = nutrient criterion for adjoining estuarine segment

$S_{TC}$  = mean salinity for tidal creek segment

$S_{FW}$  = mean salinity for adjoining/upstream freshwater segment

$S_{Est}$  = mean salinity for adjoining estuarine segment

As part of the stream assessment the physical water quality and chemical water chemistry of a stream are measured. These data only indicate a snapshot of the stream's water quality; however they are useful when compared to the trend data available from Hillsborough County Environ-

mental Protection Commission or other sources. Table 49 contains the summary water quality data and index values and adjusted values calculated from these data. The total phosphorus (TP), total nitrogen (TN) and chlorophyll a water chemistry sample data are the results of chemical analysis of samples taken during the assessment and analyzed by the Hillsborough County Environmental Protection Commission laboratory.

The growth of plants (planktonic algae, macrophytic algae and rooted plants) is directly dependent on the available nutrients within the water column of a stream and to some extent the nutrients which are held in the sediment and the vegetation biomass of a stream. Additionally, algae and other plant growth are [limited](#) by the nutrient in lowest concentration relative to that needed by a plant. Plant biomass contains less phosphorus by weight than nitrogen so phosphorus is many times the limiting nutrient. When both nutrients are present at a concentration in the stream so that either or both may restrict plant growth, the limiting factor is called “balanced”. The ratio of total nitrogen to total phosphorous, the “N to P” ratio (N/P), is used to determine the limiting factor. If N/P is greater than or equal to 30, the stream is considered phosphorus limited, when this ratio is less than or equal to 10, the stream is considered nitrogen limited and if between 10 and 30 it is considered balanced.

**Table 4. Water Quality Parameters (Laboratory) for Bullfrog Creek**

Parameter	Value	Mean Value
<b>Total Phosphorus (ug/L)</b>	<b>295.50</b>	<b>322.89</b>
<b>Total Nitrogen (ug/L)</b>	<b>1068.67</b>	<b>1182.17</b>
<b>Chlorophyll a (ug/L)</b>	<b>8.10</b>	<b>11.07</b>
<b>TN/TP</b>	<b>3.6</b>	<b>3.7</b>
<b>Limiting Nutrient</b>	<b>Nitrogen</b>	<b>Nitrogen</b>
<b>Chlorophyll TSI</b>	<b>46</b>	<b>51</b>
<b>Phosphorus TSI</b>	<b>87</b>	<b>89</b>
<b>Nitrogen TSI</b>	<b>61</b>	<b>63</b>
<b>TSI</b>	<b>53</b>	<b>57</b>
<b>Color (PCU)</b>	<b>42.30</b>	<b>49.74</b>
<b>Secchi disk depth (ft)</b>	<b>2.75</b>	<b>2.19</b>
<b>Impaired TSI for Stream</b>	<b>60</b>	<b>60</b>
<b>Stream Status (Water Column)</b>	<b>Not Impaired</b>	<b>Not Impaired</b>

The color of a stream is also important to the growth of algae. Dark, tannic streams tend to suppress algal growth and can tolerate a higher amount of nutrient in their water column; while clear streams tend to support higher algal growth with the same amount of nutrients. The color of a stream, which is measured in a unit called the “cobalt platinum unit (PCU)” because of the standard used to determine color, is important because it is used by the State of Florida to determine stream impairment as explained earlier. Rivers, streams or other “flow through” systems tend to support lower algal growth for the same amount of nutrient concentration. All these factors are important to the understanding of your stream’s overall condition. Table 9 includes many of the factors that are typically used to determine the actual state of plant growth in your stream. These data should be understood and reviewed when establishing a management plan for a stream; however, as stated above other factors must be considered when developing such a plan. Please contact the [Water Atlas Program](#) if you have questions about this part or any other part of this report.

Table 5 contains the field data taken in the upstream and downstream extents of the stream using a multi-probe (we use either a YSI 6000 or a Eureka Manta) which has the ability to directly measure the temperature, pH, dissolved oxygen (DO), percent DO (calculated from DO, temperature and conductivity). These data are listed for three levels in the stream and twice for the surface measurement. These three locations cover the predominantly freshwater portion upstream, the mixing zone and the confluence with the receiving estuary.



**Table 5. Water Chemistry Data Based on Manta Water Chemistry Probe for {LAKE\_NAME}**

Sample Location	Sample Depth (m)	Time	Temp (deg C)	Conductivity (mS/cm3)	Dissolved Oxygen (%)	Dissolved Oxygen (mg/L)	pH
Bottom - I75	0.92	6/14/2012 12:00:00 AM	26.36	0.421	77.32	6.35	6.93
Bottom - mouth	1.51	6/14/2012 12:00:00 AM	29.70	34.433	112.87	7.75	7.81
Bottom - US41	1.82	6/14/2012 12:00:00 AM	29.19	25.036	33.65	2.41	6.99
Middle - I75	0.62	6/14/2012 12:00:00 AM	26.36	0.421	77.36	6.36	6.93
Middle - mouth	0.90	6/14/2012 12:00:00 AM	29.99	33.634	117.37	8.07	7.85
Middle - US41	1.06	6/14/2012 12:00:00 AM	29.40	22.627	37.03	2.67	6.99
Surface - I75	0.44	6/14/2012 12:00:00 AM	26.37	0.421	77.37	6.36	6.93
Surface - mouth	0.44	6/14/2012 12:00:00 AM	30.26	33.200	119.68	8.21	7.84
Surface - US41	0.52	6/14/2012 12:00:00 AM	29.94	15.070	40.94	3.01	7.09

To better understand many of the terms used in this report, we recommend that the reader visit the [Hillsborough County & City of Tampa Water Atlas](#) and explore the “Learn More” areas which are found on the resource pages. Additional information can also be found using the [Digital Library](#) on the Water Atlas website.

## Section 4: Conclusion

Bullfrog Creek is a medium area (109.9-acre) stream that would be considered in the healthy category of streams based on water chemistry. It has a plant diversity of 98 species relative to the total watershed plant diversity of 98 species with about 0% percent of the open water areas containing submerged aquatic vegetation. Vegetation helps to maintain the nutrient balance in the stream as well as provide good fish habitat. The stream has many open water areas to support various types of recreation and has a fair diversity of plant species. The primary pest plants in the stream include *schinus terebinthifolius* and *leucaena leucocephala*.

This assessment was accomplished to assist stream property owners to better understand and manage their streams. Hillsborough County supports this effort as part of their [Stream Water-watch Program \(SWW\)](#) and has developed guidelines for stream property owner groups to join the SWW and receive specific assistance from the County in the management of their stream. For additional information and recent updates please visit the [Hillsborough County & City of Tampa Water Atlas](#) website.

## Stream Assessment Notes

**NOTE 1: The Water Quality Index (WQI)** is used for streams, black waters (natural tea and coffee-colored waters), and springs, while the Trophic State Index (TSI) is used for lakes and estuaries. The WQI is calculated by averaging the values of most or all of the parameters within five water quality parameter categories: 1) water clarity (measured as turbidity and/or Secchi disk depth), 2) dissolved oxygen, 3) oxygen demanding substances (measured as biochemical oxygen, chemical oxygen demand and/or total organic carbon), 4) nutrients (measured as total nitrogen, nitrite plus nitrate, and/or total phosphorus), and 5) bacteria (total coliform and-or fecal coliform).

Water Atlas presents WQIs over the last four seasons (three month intervals). The WQI "value" for a waterbody is determined by averaging the values (data) of the aforementioned parameters for each "season" (Jan-Mar, Apr-Jun, Jul-Sep, Oct-Dec). These seasonal averages are then averaged to provide an overall "rating" or WQI. The term "confidence" expresses the degree of completeness of the index; in other words, "confidence" states how many parameter categories were used to calculate the Overall Water Quality Index.

Ranges of WQI values have been established to provide a general ranking of the waterbody (Figure 1.) WQI values may also include the 'Confidence' (Figure 2), which provides you with some relative idea as to how much information was used to calculate the WQI for that waterbody.

*Note: The acronym WQI also stands for "Water Quality Inspection" in much of the DEP literature.*

WQI	Rating
0-45	Good
45-60	Fair
>60	Poor

**Figure 1. Water Quality Index (WQI) ranges and their designations.**

WQI	Rating	Confidence	Season
30	Good	5/5	Winter (2000)
40	Good	3/5	Fall (2000)
30	Good	2/5	Summer (2000)
50	Fair	3/5	Summer (2000)

**Figure 2. WQI rankings are provided with examples of Confidence values.**

**NOTE 2: Definition of a "Stream" from 62-302.531 Florida Administrative Code (FAC):**

"Stream" shall mean, for purposes of interpreting the narrative nutrient criterion in paragraph 62-302.530(47)(b), F.A.C., under paragraph 62-302.531(2)(c), F.A.C., a predominantly fresh surface waterbody with perennial flow in a defined channel with banks during typical climatic and hydrologic conditions for its region within the state. During periods of drought, portions of a stream channel may exhibit a dry bed, but wetted pools are typically still present during these conditions. Streams do not include:

non-perennial water segments where fluctuating hydrologic conditions, including periods of desiccation, typically result in the dominance of wetland and/or terrestrial taxa (and corresponding reduction in obligate fluvial or lotic taxa), wetlands, or portions of streams that exhibit lake characteristics (e.g., long water residence time, increased width, or predominance of biological taxa typically found in non-flowing conditions) or tidally influenced segments that fluctuate between predominantly marine and predominantly fresh waters during typical climatic and hydrologic conditions; or



ditches, canals and other conveyances, or segments of conveyances, that are man-made, or predominantly channelized or predominantly physically altered and;

are primarily used for water management purposes, such as flood protection, stormwater management, irrigation, or water supply; and

have marginal or poor stream habitat or habitat components, such as a lack of habitat or substrate that is biologically limited, because the conveyance has cross sections that are predominantly trapezoidal, has armored banks, or is maintained primarily for water conveyance.

**NOTE 3: The “Stream Condition Index (SCI)”** shall mean a Biological Health Assessment that measures stream biological health in predominantly freshwaters using benthic macroinvertebrates, performed and calculated using the Standard Operating Procedures for the SCI in the document titled SCI 1000: *Stream Condition Index Methods* (DEP-SOP-003/11 SCI 1000) and the methodology in *Sampling and Use of the Stream Condition Index (SCI) for Assessing Flowing Waters: A Primer* (DEP-SAS-001/11), both dated 10-24-11, which are incorporated by reference herein. Copies of the documents may be obtained from the Department’s website at <http://www.dep.state.fl.us/water/wqssp/swq-docs.htm> or by writing to the Florida Department of Environmental Protection, Standards and Assessment Section, 2600 Blair Stone Road, MS 6511, Tallahassee, FL 32399-2400. For water quality standards purposes, the Stream Condition Index shall not apply in the South Florida Nutrient Watershed Region.

**NOTE 4: Definition of a Tidal Stream:** Tidally influenced segments that fluctuate between predominantly marine and predominantly fresh waters during typical climatic and hydrologic conditions (excerpt from above FAC definitions).

For streams (other than exceptions listed above), if a site specific interpretation pursuant to paragraph 62-302.531(2)(a) or (2)(b), FAC, has not been established (see at:

<http://www.hillsborough.wateratlas.usf.edu/upload/documents/62-302.pdf>), biological information shall be used to interpret the narrative nutrient criterion in combination with Nutrient Thresholds. The narrative nutrient criterion in paragraph 62-302.530(47)(b), FAC., shall be interpreted as being achieved in a stream segment where information on chlorophyll a levels, algal mats or blooms, nuisance macrophyte growth, and changes in algal species composition indicates there are no imbalances in flora or fauna, and either:

the average score of at least two temporally independent SCIs performed at representative locations and times is 40 or higher, with neither of the two most recent SCI scores less than 35, or

the nutrient thresholds set forth in the table below are achieved.

<u>Nutrient Watershed Region</u>	<u>Total Phosphorus Nutrient Threshold<sup>1</sup></u>	<u>Total Nitrogen Nutrient Threshold<sup>1</sup></u>
<u>Panhandle West</u>	<u>0.06 mg/L</u>	<u>0.67 mg/L</u>
<u>Panhandle East</u>	<u>0.18 mg/L</u>	<u>1.03 mg/L</u>
<u>North Central</u>	<u>0.30 mg/L</u>	<u>1.87 mg/L</u>
<u>Peninsular</u>	<u>0.12 mg/L</u>	<u>1.54 mg/L</u>
<u>West Central</u>	<u>0.49 mg/L</u>	<u>1.65 mg/L</u>
<u>South Florida</u>	<u>No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.</u>	<u>No numeric nutrient threshold. The narrative criterion in paragraph 62-302.530(47)(b), F.A.C., applies.</u>

<sup>1</sup>These values are annual geometric mean concentrations not to be exceeded more than once in any three calendar year period.

**NOTE 5: Tidal Creeks** On March 15, 2013 the USEPA and the FDEP agreed that the FDEP proposed standards (62-302.532 FAC, Estuary-Specific Numeric Interpretations of the Surface Wa-

ter Quality Standards; see at: <http://www.hillsborough.wateratlas.usf.edu/upload/documents/62-302.pdf>) would be used to determine impairment in all streams. As above, this criterion allows the use of narrative standards for tidal streams but adopts those above for the majority of freshwater streams in Florida. Narrative Criteria, 62-302-530(47)(b), FAC, provides that “[i]n no case shall nutrient concentrations of a body of water be altered so as to cause an imbalance in natural populations of aquatic flora or fauna.”

Since no actual standard exists for tidal creeks we elected to use the USEPA approach that is outlined below.

***Tidal Creek Overview*** From the (Technical Support Document for U.S. EPA’s Proposed Rule for Numeric Nutrient Criteria, Volume 1 Estuaries, November 30, 2012):

“Tidal creeks and associated marshes and mangroves are refuges for small forage fish and for juveniles of larger fish to which they are considered an important spawning and nursery habitat. For example, juveniles of common snook (*Centropomus undecimalis*) depend on tidal creeks for shelter from larger predators (Adams 2005). Dominant aquatic animals in tidal creeks include mummichogs (Fundulidae) and grass shrimp (Palaeomonidae), but many other estuarine species also thrive in these habitats (Greenwood et al. 2009; Krebs et al. 2009; Janicki Environmental 2011). In general, undisturbed tidal creeks in Florida have higher fish densities than adjacent open waters. Tidal creeks can be degraded by suburban and urban development in their watersheds. Stressors from watershed development include hydrologic modification because of increased flashiness from impervious surfaces; channelization for marinas and docks; and nutrient pollution from lawn fertilizers, urban and agricultural runoff, and septic systems. As a result, tidal creeks draining developed areas have higher nutrient, chlorophyll, and fecal coliform bacteria concentrations compared to streams draining undeveloped watersheds (Holland et al. 2004; Mallin et al. 2004). Furthermore, hypoxic episodes are more extreme (prolonged and with lower dissolved oxygen) in developed watersheds than in undeveloped watersheds (Holland et al. 2004). In addition to increased nutrient concentrations, watershed development results in increased variability and volume of runoff during and after rainfall. The runoff surges cause more rapid and more extreme salinity changes as well as increased scour and changes in channel morphology. Tidal creeks with watersheds that have high impervious surface area have been observed to support degraded fish and invertebrate communities in South Carolina. Although commercially important spot and shrimp populations were reduced in affected creeks, mummichog and grass shrimp remained (Holland et al. 2004; Lerberg et al. 2000; Mallin et al. 2004). Other studies have shown that low-salinity waters of tidal creeks in developed areas can develop nuisance algal bloom conditions (Mallin et al. 2004; MacPherson et al. 2007), with the bloom waters moving back and forth with the tides. Such bloom conditions can also contribute to more severe hypoxic episodes.

***Derivation of Numeric Nutrient Criteria for Tidal Creeks:*** Tidal creeks were classified separately from estuaries because tidal creeks are expected to have higher nutrient and chlorophyll concentrations than adjacent, open waters. The classification and segmentation approach used for estuaries was not considered practical because of the large number and variety of small systems. A definitional approach was chosen, applicable to all tidal creeks, to be implemented on a case-by-case basis as data allow. Several options were considered for deriving numeric nutrient criteria for tidal creeks, including applying inland freshwater criteria derived for upstream waters or applying estuarine criteria derived for downstream waters. Neither of those two approaches alone would be applicable to the full range and variability of tidal creeks. Ultimately, EPA selected two approaches for deriving numeric TN and TP criteria that account for the inherent variability of tidal creeks.

*The first approach* is to apply separately derived inland TN and TP criteria for adjacent freshwaters if the mean chloride of the tidal creek is less than 1,500 mg/L, or apply estuarine TN and TP criteria for adjacent downstream waters if the mean chloride of the tidal creek is greater than or equal to 1,500 mg/L.

*The second approach* uses linear interpolation to derive criteria for TN and TP for tidal creeks using criteria that were derived separately for adjacent inland freshwater and estuary areas on

the basis of mean salinity. Criteria would be derived by that method only where there are sufficient salinity data to allow for interpolation. The calculation uses the following formula:

$$C_{TC} = C_{FW} + (S_{TC} - S_{FW}) \times \left( \frac{C_{Est} - C_{FW}}{S_{Est} - S_{FW}} \right)$$

where

$C_{TC}$  = nutrient criterion for tidal creek segment

$C_{FW}$  = nutrient criterion for adjoining/upstream freshwater segment

$C_{Est}$  = nutrient criterion for adjoining estuarine segment

$S_{TC}$  = mean salinity for tidal creek segment

$S_{FW}$  = mean salinity for adjoining/upstream freshwater segment

$S_{Est}$  = mean salinity for adjoining estuarine segment

Example:

Segment	Mean Salinity (ppt)	Criterion Concentration
Freshwater segment	0.5	2.5
Tidal creek segment	20	$C_{TC}$
Estuarine segment	30	0.8

$$C_{TC} = 2.5 + (20 - 0.5) \times \left( \frac{0.8 - 2.5}{30 - 0.5} \right) = 1.376$$

**Tampa Bay Tidal Creeks** (From Letter Memorandum, Titled *Tampa Bay Numeric Nutrient Criteria: Tidal Creeks*, prepared by Janicki Environmental, Inc. for Tampa Bay Estuary Program. 16 February 2011):

“There are approximately sixty tidal creeks that are terminal tributaries to Tampa Bay or to smaller embayments within the bay (Figure 3). Tampa Bay tidal creeks differ substantially in scale from the larger tidal rivers and these differences in relative channel geomorphology result in disparate hydrological and physicochemical characteristics from Tampa Bay’s tidal rivers. Some of the larger tidal creeks extend far enough into the watershed that they have lower order, freshwater tributaries that feed into them (e.g., Bullfrog Creek, Double Branch Creek, Frog Creek). Tidal creeks also differ from freshwater tributaries of the same size primarily due to their connection to the estuary. Small freshwater tributaries do not experience the semidiurnal tides which cause the daily and even hourly fluctuations in water level, flow direction, salinity, water temperature and dissolved oxygen (DO) often recorded in tidal creeks (Buzzelli et al., 2007). Delineation of estuarine and freshwater tributaries to Tampa Bay is provided in Figure 3 below. Unmodified tidal creeks are characterized by sinuous, meandering channels with average water depths <1.0 m, while those creeks modified for drainage, mosquito control, or navigation often have straightened channels with steeper, more uniform banks than unmodified creeks. Tidal creeks altered for navigation are typically deeper than other creeks (>2.0 m in depth) and often have hardened shorelines that have been cleared of vegetation. Most tidal creeks in Tampa Bay are relatively narrow, spanning only 25-50 m from bank to bank, in contrast to the tidal rivers which are 100-300 m wide



on average, although some of the larger tidal creeks reach 100 m or more in width near the mouth. The bathymetry of tidal creeks consists of alternating areas of deep, erosional and shallow, depositional bottom, unless the creek has been channelized, in which case, it is often uniformly deep.”



**Figure 3. Named Tidal Creeks in Tampa Bay Region. From Letter Memorandum, Titled Tampa Bay Numeric Nutrient Criteria: Tidal Creeks, prepared by Janicki Environmental, Inc. for Tampa Bay Estuary Program. 16 February 2011.**

Tidal Creeks of interest for our reports include those that flow to Old Tampa Bay, Hillsborough Bay and Middle Tampa Bay. In our assessment we will then use the freshwater stream criteria for tidal creeks segments if a chloride concentration less than or equal to 1,500 mg/L and estuary criteria (see below) for segments with a chloride concentration of greater than 1,500 mg/L.

(1) Estuary-specific numeric interpretations of the narrative nutrient criterion in paragraph 62-302.530(47)(b), FAC, are in the table below. The concentration-based estuary interpretations are open water, area-wide averages. The interpretations expressed as load per million cubic meters of freshwater inflow are the total load of that nutrient to the estuary divided by the total volume of freshwater inflow to that estuary.

Estuary	Total Phosphorus	Total Nitrogen	Chlorophyll <i>a</i>
(a) Clearwater Harbor/St. Joseph Sound	Annual geometric mean values not to be exceeded more than once in a three year period. Nutrient and nutrient response values do not apply to tidally influenced areas that fluctuate between predominantly marine and predominantly fresh waters during typical climatic and hydrologic conditions.		
1. St Joseph Sound	0.05 mg/L	0.66 mg/L	3.1 µg/L
2. Clearwater North	0.05 mg/L	0.61 mg/L	5.4 µg/L
3. Clearwater South	0.06 mg/L	0.58 mg/L	7.6 µg/L
(b) Tampa Bay	Annual totals for nutrients and annual arithmetic means for chlorophyll <i>a</i> , not to be exceeded more than once in a three year period. Nutrient and nutrient response values do not apply to tidally influenced areas that fluctuate between predominantly marine and predominantly fresh waters during typical climatic and hydrologic conditions.		
1. Old Tampa Bay	0.23 tons/million cubic meters of water	1.08 tons/million cubic meters of water	9.3 µg/L
2. Hillsborough Bay	1.28 tons/million cubic meters of water	1.62 tons/million cubic meters of water	15.0 µg/L
3. Middle Tampa Bay	0.24 tons/million cubic meters of water	1.24 tons/million cubic meters of water	8.5 µg/L
4. Lower Tampa Bay	0.14 tons/million cubic meters of	0.97 tons/million cubic meters of	5.1 µg/L

Note: 1 ton / (million cubic meter) = 0.907 mg/L

**NOTE 6: Salinity** Salinity is a way of expressing the “saltiness” or dissolved salt content (primarily sodium chloride, magnesium and calcium sulfates and bicarbonates) of natural waters and is normally only used for saltwater systems. The unit of salinity commonly used is a part per thousand (ppt). Natural water salinity regimes commonly discussed in the literature include freshwater (< 0.05 ppt), Oligohaline (0.05-0.5 ppt), mesohaline (0.5-5 ppt), polyhaline (5-18 ppt), mixoeuhaline (18-30 ppt) and metahaline (30-40) ppt. Seawater in the open ocean is normally in the metahaline regime.

The salinity of a natural water is an important factor to measure and to understand. Salinity can be used to trace the movement of estuarine waters within a tidal stream which is a factor of tide and wind velocity. It is also important in understanding the types of organisms that might be expected to exist in a specific segment of a tidal stream. Additionally, salinity influences the kinds of [plants](#) that will grow either in a in the stream or along the wetland margin of a stream. A plant adapted to saline conditions is called a [halophyte](#). Organisms (mostly bacteria) that can live in very salty conditions are classified as [extremophiles](#), or [halophiles](#) specifically. An organism that can withstand a wide range of salinities is [euryhaline](#).

One of the criteria that has been proposed by the USEPA in their technical volume on estuary and tidal creek numeric nutrient criteria is based on the chloride concentration for a stream. They propose a chloride concentration of 1,500 mg/L as the point where a stream should be classified as tidal. That is a stream segment with a chloride concentration greater that this value is a tidal segment. Looking at the table below which gives a relationship between salinity, conductivity and chloride concentration, this value converts to a salinity of 2.47 ppt.

**Relationship between Electrical Conductivity (EC -  $\mu\text{S}/\text{cm}$ ), salinity concentration, chloride concentration and Practical Salinity Units**


EC @25°C	Salinity ppt or g/l	Salinity ppm or mg/l	Cl ppm or mg/l	PSU	EC @25°C	Salinity ppt or g/l	Salinity ppm or mg/l	Cl ppm or mg/l	PSU
100	0.05	51	31	0.07	5000	2.68	2678	1626	2.41
200	0.10	97	59	0.12	6000	3.25	3253	1975	2.88
300	0.14	144	88	0.17	7000	3.84	3836	2328	3.33
400	0.19	193	117	0.22	8000	4.43	4426	2687	3.79
500	0.24	241	147	0.27	9000	5.02	5022	3049	4.24
600	0.29	291	177	0.32	10000	5.62	5625	3414	4.69
700	0.34	341	207	0.37	12000	6.85	6846	4155	5.57
800	0.39	391	237	0.42	14000	8.09	8087	4909	6.44
900	0.44	442	268	0.47	16000	9.35	9345	5673	7.31
1000	0.49	493	299	0.52	18000	10.62	10621	6447	8.15
1200	0.60	596	362	0.61	20000	11.91	11911	7230	8.99
1400	0.70	699	425	0.71	22000	13.22	13215	8022	9.82
1600	0.80	804	488	0.81	24000	14.53	14534	8822	10.64
1800	0.91	910	552	0.90	26000	15.86	15865	9630	11.45
2000	1.02	1016	617	1.00	28000	17.21	17208	10446	12.25
2200	1.12	1123	682	1.10	30000	18.56	18564	11268	13.04
2400	1.23	1231	747	1.19	32000	19.93	19931	12098	13.82
2600	1.34	1339	813	1.29	34000	21.31	21309	12935	14.60
2800	1.45	1448	879	1.38	36000	22.70	22699	13778	15.37
3000	1.56	1558	946	1.48	38000	24.10	24099	14629	16.13
3200	1.67	1668	1012	1.57	40000	25.51	25510	15485	16.88
3400	1.78	1778	1080	1.67	42000	26.93	26932	16348	17.63
3600	1.89	1889	1147	1.76	44000	28.36	28364	17217	18.37
3800	2.00	2001	1215	1.85	46000	29.81	29806	18093	19.10
4000	2.11	2113	1283	1.95	48000	31.26	31259	18975	19.83
4200	2.23	2225	1351	2.04	50000	32.72	32722	19863	20.56
4400	2.34	2338	1419	2.13	52000	34.19	34195	20757	21.27
4600	2.45	2451	1488	2.23	54000	35.68	35678	21657	21.99
4800	2.56	2565	1557	2.32					

**NB There is no direct conversion between many of these units.**

**The above information is a guide only and where relationships between values have been established, they are specific to Hickling Broad water quality data only**

**Vegetation Zones:** The three primary aquatic vegetation zones are shown below:





An **adjusted chlorophyll a value** ( $\mu\text{g/L}$ ) was calculated by modifying the methods of Canfield et al (1983). The total wet weight of plants in the stream (kg) was calculated by multiplying stream surface area ( $\text{m}^2$ ) by PAC (percent area coverage of macrophytes) and multiplying the product by the biomass of submersed plants ( $\text{kg wet weight m}^2$ ) and then by 0.25, the conversion for the 1/4 meter sample cube. The dry weight (kg) of plant material was calculated by multiplying the wet weight of plant material (kg) by 0.08, a factor that represents the average percent dry weight of submersed plants (Canfield and Hoyer, 1992) and then converting to grams. The potential phosphorus concentration ( $\text{mg/m}^3$ ) was calculated by multiplying dry weight (g) by 1.41 mg TP g<sup>-1</sup> dry weight, a number that represents the mean phosphorus (mg) content of dried plant material measured in 750 samples from 60 Florida lakes (University of Florida, unpublished data), and then dividing by stream segment volume ( $\text{m}^3$ ) and then converting to  $\mu\text{g/L}$  (1000/1000). From the potential phosphorus concentration, a predicted chlorophyll a concentration was determined from the total phosphorus and chlorophyll a relationship reported by Brown (1997) for 209 Florida lakes. Adjusted chlorophyll a concentrations were then calculated by adding each lake's measured chlorophyll a concentration to the predicted chlorophyll a concentration.

**Wide Area Augmentation System (WAAS)** is a form of differential GPS (DGPS) where data from 25 ground reference stations located in the United States receive GPS signals from GPS satellites in view and retransmit these data to a master control site and then to geostationary satellites. The geostationary satellites broadcast the information to all WAAS-capable GPS receivers. The receiver decodes the signal to provide real time correction of raw GPS satellite signals also received by the unit. WAAS-enabled GPS is not as accurate as standard DGPS which employs close by ground stations for correction, however; it was shown to be a good substitute when used for this type of mapping application. Data comparisons were conducted with both types of DGPS employed simultaneously and the positional difference was determined to be well within the tolerance established for the project.