

# LAKE ASSESSMENT REPORT FOR LAKE MAGDALENE IN HILLSBOROUGH COUNTY FLORIDA

Date Assessed: 6/15/2006

Assessed by: Sarah Koenig and David Eilers

Reviewed by: Jim Griffin, Ph.D.

## **INTRODUCTION**

This assessment was conducted to update existing physical and ecological data for Lake Magdalene (lake view see Figure 1) on the Hillsborough County Watershed Atlas (<http://www.hillsborough.wateratlas.usf.edu/>). The project is a collaborative effort between the University of South Florida's Center for Community Design and Research and Hillsborough County Stormwater Management Section. The project is funded by Hillsborough County and the Southwest Florida Water Management District's Northwest Hillsborough, Hillsborough River and Alafia River Basin Boards. The project has, as its primary goal, the rapid assessing of up to 150 lakes in Hillsborough County during a five year period. The product of these investigations will provide the County, lake property owners and the general public a better understanding of the general health of Hillsborough County lakes, in terms of shoreline development, water quality, lake morphology (bottom contour, volume, area etc.) and the plant biomass and species diversity. These data are intended to assist the County and its citizens to better manage lakes and lake centered watersheds.

**Figure 1. General Photo of Lake Magdalene.**



**The first section** of the report provides the results of the overall morphological assessment of the lake. Primary data products include: a contour (bathymetric) map of the lake, area, volume and depth statistics, and the water level at the time of assessment. These data are useful for evaluating trends and for developing management actions such as plant management where depth and lake volume are needed.

**The second section** provides the results of the vegetation assessment conducted on the lake. These results can be used to better understand and manage vegetation in your lake. A list is provided with the different plant species found at various sites around the lake. Potentially invasive, exotic (non-native) species are identified in a plant list and the total non-native is presented in a summary table. Watershed values provide a means of reference.

**The third section** provides the results of the water quality sampling of the lake. Both field data and laboratory data are presented. The trophic state index (TSI)<sup>i</sup> is used to develop a general lake health statement, which is calculated for both the water column with vegetation and the water column if vegetation were removed. These data are derived from the water chemistry and vegetative submerged biomass assessments and are useful in understanding the results of certain lake vegetation management practices.

The intent of this assessment is to provide a starting point from which to track changes in your lake, and where previous comprehensive assessment data is available, to track changes in the lake's general health. These data can provide the information needed to determine changes and to monitor trends in physical condition and ecological health of the lake.

## Section 1: Lake Morphology

**Bathymetric Map<sup>ii</sup>**. The bottom of the lake was mapped using a Lowrance LCX 26C HD Wide Area Augmentation System (WAAS)<sup>iii</sup> enabled Global Positioning System (WAAS-GPS) with fathometer (bottom sounder) to determine the boat's position, and bottom depth in a single measurement. The result is an estimate of the lake's area, mean and maximum depths, and volume and the creation of a bottom contour map (Figure 2). Besides pointing out the deeper fishing holes in the lake, the morphologic data derived from this part of the assessment can be valuable to overall management of the lake vegetation as well as providing flood storage data for flood models. Table 1 provides the lake's morphologic parameters in various units.

**Table 1. Lake Area Depth and Volume**

Parameter	Feet	Meters	Acres	Gallons
Surface Area (sq)	10,445,932.86	970,470.47	239.81	
Mean Depth	8.10			
Maximum Depth	22.00			
Volume (cubic)	84,253,114.23	2,385,782.80		630,257,251.82
Gage (above datum)	49.05			




Figure 1. Contour map for Lake Magdalene. The lake was not remapped in 2006 because the previous map shown here (2003) was constructed within the last 5 years. The mapping technique used in 2006 employs a standard DGPS for horizontal position and a fathometer for depth.



## Section 2: Lake Ecology (vegetation)

The lake's apparent vegetative cover and shoreline detail are evaluated using the aerial shown in Figure 3 and by use of GPS. Submerged vegetation is determined from evenly spaced contours sampled using a Lowrance 26c HD, combined GPS/fathometer described earlier. Twenty vegetation assessment sites were used for Lake Magdalene (Figure 3). Fifteen sites for intensive sampling as dictated by the *Lake Assessment Protocol* (copy available on request) for a lake of this size. Five sites were used to *ground truth* the apparent vegetative cover for vegetated islands shown in aerial photographs. The site positions are set using GPS and then loaded into a GIS mapping program (ArcGIS) for display. Each site is sampled in the three primary vegetative zones (emergent, submerged and floating)<sup>iv</sup>. The latest aeriels (2005, 6 inch resolution, SWFWMD aeriels) are used to provide shore details (docks, structures, vegetation zones) and to calculate the extent of surface vegetation coverage. The primary indices of submerged vegetation cover and biomass for the lake, percent area coverage (PAC) and percent volume infestation (PVI), are determined by transiting the lake by boat and employing a fathometer to collect "hard and soft return" data. These data are later analyzed for presence and absence of vegetation and to determine the height of vegetation if present. The PAC index is determined from the presence and absence analysis of 100 sites in the lake and the PVI index is determined by measuring the difference between hard returns (lake bottom) and soft returns (top of vegetation) for sites (within the 100 analyzed sites) where plants are determined present.

The data collected during the site vegetation sampling include vegetation type, exotic vegetation, predominant plant species and submerged vegetation biomass. The total number of species from all sites is used to approximate the total diversity of aquatic plants and the total non-native plants on the lake (Table 2). The Watershed value in Table 2 only includes lakes sampled during the lake assessment project begun in May of 2006. These data will change as additional lakes are sampled. Tables 3 through 7 detail the results from the 2006 aquatic plant assessment for you lake. These data are determined from the 15 sites used for intensive vegetation surveys.

The tables are divided into Floating Leaf, Emergent and Submerged plants and contain the plant code, species, common name and presence (1) or absence (blank) of species and the calculated percent occurrence (number sites species is found/number of sites) and type of plant (Native, Non-Native, Invasive, Pest). In the "Type" category, the term invasive indicates the plant is commonly considered invasive in this region of Florida and the term "Pest" indicates that the plant has a greater than 55% occurrence in your lake and is also considered a problem plant for this region of Florida. These two terms are somewhat subjective; however, they are provided to give lake property owners some guidance in the management of plants on their property. Please remember that to remove or control plants in a wetland (lake shoreline) in Hillsborough County the property owner must secure an [Application To Perform Miscellaneous Activities In Wetlands](http://www.epchc.org/forms_documents.htm) ([http://www.epchc.org/forms\\_documents.htm](http://www.epchc.org/forms_documents.htm)) permit from the Environmental Protection Commission of Hillsborough and for management of in-lake vegetation outside the wetland fringe (for lakes with an area greater than 10 acres), the property owner must secure a Florida Department of Environmental Protection permit (<http://www.dep.state.fl.us/lands/invaspec/>).

**Table 2 Total diversity, percent exotics, and number of EPPC pest plants**

Parameter	Lake	Watershed
Total Plant Diversity (# of Taxa)	54	98
Total Non-Native Plants	9	17
Total Pest Plant Species	5	17



Figure 2. 2004 six inch resolution aerial and vegetation assessment sites on Lake Magdalene.



**Table 3. List of Floating Leaf Zone Aquatic Plants Found**

Code	Plant Species	Common Name	Sample Sites															% Occurrence	Native, Non-Native (NN), Invasive (I), Pest (P)
			1	2	3	4	5	6	7	8	9	10	11	12	13	14	15		
SMA	<i>Salvinia minima</i>	Water Spangles, Water Fern							1				1					13.33%	NN-I
MMA	<i>Marsilea mutica</i>	Water Clover								1								6.67%	NN-I
NOA	<i>Nymphaea odorata</i>	American White Water lily, Fragrant Water Lily	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	100%	Native
NLM	<i>Nuphar lutea</i> var. <i>advena</i>	Spatterdock, Yellow Pondlily		1	1	1			1	1	1		1			1	1	60%	Native
HYE	<i>Hydrocotyl umbellata</i>	Manyflower Marshpennywort, Water Pennywort		1			1	1	1	1					1	1		46.67%	Native



**Table 4 List of Emergent Zone Aquatic Plants Found**

Code	Plant Species	Common Name	Sample Sites															% Occurrence	Native (N), Non-Native (NN), Invasive (I), Pest (P)	
			1	2	3	4	5	6	7	8	9	10	11	12	13	14	15			
MEL	Melaleuca quinquenervia	Punk Tree, Melaleuca		1	1	1	1			1		1	1		1	1		1	66.67%	NN – I - P
APS	Alternanthera philoxeroides	Alligator Weed	1	1			1									1			26.67%	NN- I
CCA	Cinnamomum camphora	Camphor-tree								1		1	1						20%	NN - I
PRS	Panicum repens	Torpedo Grass	1				1										1		20%	NN - I
WTA	Sphagneticola (Wedelia) trilobata	Creeping Oxeye					1			1							1		20%	I
CLA	Casuarina equisetifolia	Australian Pine								1									6.67%	I
FSC	Fuirena spp.	Rush Fuirena	1	1	1	1	1	1	1	1				1	1	1	1	1	80%	Native
SLA	Sagittaria lancifolia	Bulltongue Arrowhead, Duck Potato			1	1	1	1	1	1				1	1	1	1	1	66.67%	Native
TYP	Typha spp.	Cattails		1	1	1	1		1			1	1	1	1		1	1	66.67%	Native – I - P
PAN	Panicum spp.	Panic grasses				1	1	1		1	1	1		1	1	1			60%	Native - P
PCA	Pontederia cordata	Pickerel Weed		1	1	1		1	1					1	1		1		53.33%	Native
BCA	Bacopa caroliniana	Lemon Bacopa	1	1			1	1				1		1		1			46.67%	Native
LOP	Ludwigia spp.	Water Primroses, Primrosewillow		1	1		1	1			1	1	1						46.67%	Native
MSS	Mikania scandens	Climbing Hempvine		1		1	1	1								1	1		40%	Native
PHN	Panicum hemitomon	Maidencane				1	1	1		1	1			1					40%	Native
SAL	Salix spp.	Willow		1		1		1	1			1				1			40%	Native
ELC	Eleocharis cellulosa	Club Rush			1	1	1		1									1	33.33%	Native

**Table 5 List of Emergent Zone Aquatic Plants Found**

Code	Plant Species	Common Name	Sample Sites															% Occurrence	Native (N), Non-Native (NN), Invasive (I), Pest (P)		
			1	2	3	4	5	6	7	8	9	10	11	12	13	14	15				
BMI	Bacopa monnieri	Common Bacopa, Herb-Of-Grace					1	1						1				1		26.67%	Native
WAX	Myrica cerifera	Wax Myrtle					1						1	1				1		26.67%	Native
BOC	Boehmeria cylindrica	Bog Hemp, False Nettle		1									1					1		20%	Native
EBI	Eleocharis baldwinii	Baldwin's Spikerush, Roadgrass					1	1										1		20%	Native
POL	Polygonum spp.	Smartweed, Knotweed					1											1	1	20%	Native
SCS	Scirpus cubensis	Burhead Sedge, Cuban Scirpus					1							1				1		20%	Native
TAS	Taxodium ascendens	Pond Cypress						1		1								1		20%	Native
UNK	UNKNOWN SPP	Unidentified Plant Species	1							1				1						20%	Native
ACE	Acer rubrum var. trilobum	Southern Red Maple								1			1							13.33%	Native
COS	Cephalanthus occidentalis	Common Buttonbush								1	1									13.33%	Native
CYO	Cyperus odoratus	Fragrant Flatsedge					1					1								13.33%	Native
HTM	Hypericum tetrapetalum	Fourpetal St. John's-Wort					1											1		13.33%	Native
SCI	Scirpus spp.	Sedge											1	1						13.33%	Native
DVA	Diodia virginiana	Buttonweed						1												6.67%	Native
EAA	Eclipta alba (prostrata)	False Daisy, Yerba De Tajo					1													6.67%	Native



**Table 6 List of Emergent Zone Aquatic Plants Found**

Code	Plant Species	Common Name	Sample Sites															% Occurrence	Native (N), Non-Native (NN), Invasive (I), Pest (P)	
			1	2	3	4	5	6	7	8	9	10	11	12	13	14	15			
IRI	Iris spp.	Flag						1											6.67%	Native
MVA	Magnolia virginiana	Sweetbay Magnolia												1					6.67%	Native
OCA	Osmunda cinnamomea	Cinnamon Fern										1							6.67%	Native
PIN	Pinus spp.	Pine Tree							1										6.67%	Native
SLT	Sagittaria latifolia	Wapato, Common Arrowhead, Duck Potato					1												6.67%	Native
SAM	Sambucus canadensis	Elderberry											1						6.67%	Native



**Figure 5. Cattail Fringe in Lake Magdalene**



**Figure 6. Rush Fuirena and Melaleuca in Lake Magdalene**

**Table 7 List of Submerged Zone Aquatic Plants Found**

Code	Plant Species	Common Name	Sample Sites														% Occurrence	Comment
			1	2	3	4	5	6	7	8	9	10	11	12	13	14		
HVA	Hydrilla verticillata	Hydrilla, water thyme	1	1	1	1			1	1	1	1	1	1	1	1	86.67%	NN-I-P
NIT	Nitella spp.	Nitella						1		1		1			1		26.67%	Native
UTA	Utricularia spp.	Bladderwort	1	1	1	1	1	1	1	1	1	1	1	1	1	1	100%	Native
PIS	Potamogeton illinoensis	Pond Weed	1	1	1	1	1	1	1	1	1		1	1	1	1	93.33%	Native-I-P
CHA	Chara spp.	Muskgrass	1		1	1	1			1		1					40%	Native
POT	Potamogeton spp.	Pond Weed	1					1	1	1		1	1				40%	Native
NGS	Najas guadelupensis	Southern Waternymph		1	1				1			1		1			33.33%	Native
ALG	Algal Spp.	Algal Mats, Floating			1												6.67%	Native
CDM	Ceratophyllum demersum	Coontail											1				6.67%	Native
MGM	Micranthemum glomeratum	Manatee Mudflower, Baby's Tears													1		6.67%	Native
VAA	Vallisneria americana	Tapegrass	1														6.67%	Native



**Figure 7. Hydrilla and Illinois Pond Weed in Lake Magdalene**




**Figure 8. Floating Algae in Lake Magdalene**

### Section 3: Lake Water Chemistry

A critical element in any lake assessment is the long-term water chemistry data set. The primary source of water quality trend data for Florida lakes is the Florida LAKEWATCH volunteer and the Florida LAKEWATCH water chemistry data. Hillsborough County is fortunate to have a large cadre of volunteers who have collected lake water samples for significant time period. These data are displayed and analyzed on the Water Atlas as shown in Figure 10 for Lake Magdalene. Additional data, when available, is also included on the Water Atlas; however, the LAKEWATCH data remains the primary source. By the trend data shown in Figure 10, the lake may be considered in reasonable health in terms of the trophic state index. Lake Magdalene is a clear water lake and as such it must maintain a TSI of below 40 to not be considered impaired by the State of Florida guidelines<sup>v</sup>. Lake Magdalene's long term water quality data indicates enough violations of these criteria to be classified by Florida DEP as impaired. The more recent data indicate an improving trend and if this continues the lake may be removed from the impaired waters list. The general trend seen in Figure 10 may indicate a relationship between periods of heavy rainfall and increased TSI.

**Figure 9. Recent (through March 2006) Trophic State Index graph from Hillsborough Watershed Atlas**  
<http://www.hillsborough.wateratlas.usf.edu/lake/waterquality.asp?wbodyid=5168&wbodyatlas=lake>





As part of the lake assessment the physical water quality and chemical water chemistry of a lake are measured. These data only indicate a snap shot of the lakes water quality; however they are useful to comparing to the trend data. Table 8A contains the summary water quality data and index values and adjusted values calculated from these data. The total phosphorus (TP), total nitrogen (TN) and chlorophyll (a) water chemistry sample data are the results of chemical analysis of samples taken during the assessment and analyzed by the Hillsborough County Environmental Protection Commission laboratory. These data compare well with the mean data from the LAKEWATCH data set for the lake. The trophic state index (TSI) calculated from the sample data (37.25) is well within the values shown in Figure 10. Table 8B contains the field data taken in the center of the lake using a YSI Corporation – 6000 multi-probe which has the ability to directly measure the temperature, pH, dissolve oxygen (DO), percent DO (calculated from DO, temperature and conductivity) and Turbidity. These data are listed for three levels in the lake and twice for the surface measurement. The duplicate surface measurement was taken as a quality assurance check on measured data.

**Table 8A. Water Quality Parameters (Laboratory)**

Summary Table for Water Quality		
Parameter	Value	Comment
TP ug/L	14.00	
TN mg/L	1.07	
Chla ug/L	3.80	
Chla TSI	36.02	
TP TSI	38.48	
TN TSI	61.05	
Secchi Disk (SD)	7.22	
TSI	37.25	P limited
PAC	76%	
PVI	47%	
Adj TP	4.89	P from Veg Added
Adj TSI	40.79	With additional P

**Table 8B. Water Quality Parameters (Field-YSI)**

Sample Location	Time	Temp (°C)	Conductivity (mS/cm3)	Dissolved Oxygen (%)	DO (mg/L)	PH (SU)	ORP (ORP)	Turbidity (NTU)	Secchi Depth (ft)
Surface	13:22	29.17	0.242	109.1	8.36	8.3	254.9	0.6	
Mid	13:22	29.02	0.242	110.8	8.51	8.44	253.4	0.7	
Bottom	13:22	27.52	0.241	96.6	7.65	8.09	258.7	6	
Surface	13:22	29.19	0.242	106.4	8.16	8.35	249.7	0.7	
<b>Mean Value</b>	<b>13:22</b>	<b>28.73</b>	<b>0.242</b>	<b>105.7</b>	<b>8.17</b>	<b>8.30</b>	<b>254.2</b>	<b>6.5</b>	<b>7.22</b>

Table 8A also provides data derived from the vegetation assessment which is used to determine an adjusted TSI. This is accomplished by calculating the amount of phosphorus that could be released by existing submerged vegetation if this vegetation were treated with an herbicide or managed by the addition of Triploid grass carp (*Ctenopharyngodon idella*). While it would not be expected that all the vegetation would be turned into available phosphorus by these management methods, the data is useful when planning various management activities. Approximately 76 % of the lake has submerged vegetation present and this vegetation represents about 47 % of the available lake volume. The vegetation holds enough phosphorus to add about 4.9µg/L of the nutrient to the water column. Because the growth of algae in the water is regulated by the

availability of phosphorus (the lake is phosphorus limited), the release of this phosphorus would stimulate algal growth. These changes in the water chemistry and biology would be indicated by an increased TSI from 37.25 to about 40.8. The lake water clarity which is indicated by the Secchi Disk (SD) value at 7.22 feet would be reduced under these conditions.

#### **Section 4: Conclusion**

Lake Magdalene is a medium to large sized (240 acre) lake that would be considered in the mesotrophic (good) category of lakes based on water chemistry. It has a higher than normal concentration of aquatic vegetation and one of the few lakes in Hillsborough County with vegetated islands. About 76% of the open water areas contain submerged vegetation and this vegetation helps to maintain the nutrient balance in the lake as well as provide good fish habitat. The lake has many open water areas that support various types of recreation and has a good diversity of plant species. The primary Pest plants in the lake include Punk tree (*Melaleuca*), Hydrilla, and Pond Weed (*Potamogeton illinoensis*). For more information and recent updates please see the Hillsborough Watershed Atlas (water atlas) website at: <http://www.hillsborough.wateratlas.usf.edu/lake/waterquality.asp?wbodyid=5168&wbodyatlas=lake>


i "Trophic" means "relating to nutrition." The Trophic State Index (TSI) takes into account chlorophyll, nitrogen, and phosphorus, which are nutrients required by plant life. For more information please see *learn more* at:

<http://www.hillsborough.wateratlas.usf.edu/lake/default.asp?wbodyid=5168&wbodyatlas=lake>

ii A bathymetric map is a map that accurately depicts all of the various depths of a water body. An accurate bathymetric map is important for effective herbicide application and can be an important tool when deciding which form of management is most appropriate for a water body. Lake volumes, hydrolic retention time and carrying capacity are important parts of lake management that require the use of a bathymetric map.

iii WAAS is a form of differential GPS (DGPS) where data from 25 ground reference stations located in the United States receive GPS signals from GPS satellites in view and retransmit these data to a master control site and then to geostationary satellites. The geostationary satellites broadcast the information to all WAAS-capable GPS receivers. The receiver decodes the signal to provide real time correction of raw GPS satellite signals also received by the unit. WAAS enabled GPS is not as accurate as standard DGPS which employs close by ground stations for correction, however; it was shown to be a good substitute when used for this type of mapping application. Data comparisons were conducted with both types of DGPS employed simultaneously and the positional difference was determined to be well within the tolerance established for the project.

iv The tree primary aquatic vegetation zones are shown below:



v A lake is impaired if " (2) For lakes with a mean color less than or equal to 40 platinum cobalt units, the annual mean TSI for the lake exceeds 40, unless paleolimnological information indicates the lake was naturally greater than 40, or For any lake, data indicate that annual mean TSIs have increased over the assessment period, as indicated by a positive slope in the means plotted versus time, or the annual mean TSI has increased by more than 10 units over historical values. When evaluating the slope of mean TSIs over time, the Department shall use a Mann's one-sided, upper-tail test for trend, as described in Nonparametric Statistical Methods by M. Hollander and D. Wolfe (1999 ed.), pages 376 and 724 (which are incorporated by reference), with a 95% confidence level."

---

Excerpt from Impaired Water Rule (IWR). Please see:  
<http://www.dep.state.fl.us/water/tmdl/docs/AmendedIWR.pdf>