



The lake assessments are created in partnership with Hillsborough County and the Florida Center for Community Design and Research  
**LAKE ASSESSMENT DOCUMENT**

Lake Keene    10/14/98    Watershed: Cypress Creek

Lake assessments are being conducted to contribute physical and ecological data to the Atlas as a collaborative effort between project partners. The goal is to rapidly assess many of the lakes in the county and thus provide stakeholders a better understanding of the character of the lake, its shore, and the aquatic plants present there. These data are intended to assist in the future management of the lake and its watershed.

The first section of the report provides the results of the bottom mapping effort: a contour (bathymetric) map of the lake, area, volume and depth statistics, and the water level at the time of assessment (if available).

The second section provides the results of the ecological (vegetation) assessment conducted on the lake. These results can be used to better manage vegetation in your lake. A list is provided with the different plant species found at various sites around the lake. Potentially invasive, exotic (non-native) species are identified in a plant list and the percent of exotics is presented in a summary table. The results of this study are compared with other lakes in the watershed.

The intent of the assessment is to provide a starting point from which to track changes in your lake. These data can provide the information needed to determine changes and to monitor trends in physical condition and ecological health of the lake.

**I. Physical Data – Area, Depth, Volume, & Bottom Contours**



The bottom of the lake was mapped using a sophisticated Global Positioning System (GPS) to determine the boat’s position, and a depth-finder to provide depth associated with that measured position. The result is an estimate of your lake’s area, mean and maximum depths, and volume (Table 1) and the creation of a bottom contour map.

Table 1. Physical Characteristics of Your Lake.

Surface Area (acres):	<u>30</u>
Mean Depth (feet):	<u>8.5</u>
Maximum Depth (feet):	<u>24.2</u>
Volume (gallons):	<u>83,803,832</u>

# Lake Keene

Section - Township - Range  
7-27-19

-  Contour Lines Expressed in 2-Foot Intervals
-  Lake Perimeter Ground Level

## EXPLANATION:

Survey Date October 21, 1998


Lake water level was 62.5 ft above sea level when the lake was surveyed. Contours are expressed in absolute depth below this level.

## DATA SOURCES:

Digital orthophotos by United States Geological Survey. All contours generated by Florida Center for Community Design and Research based on survey data provided by the Hillsborough County Lake Management Program.



Hillsborough County





The lake assessments are created in partnership with Hillsborough County and the Florida Center for Community Design and Research  
**LAKE ASSESSMENT DOCUMENT**

Lake Keene 10/14/98 Watershed: Cypress Creek

**II. Ecological Data**

Aquatic Plant Survey

Approximately equispaced sites are haphazardly mapped around the lake and the aquatic plants at each site are surveyed. The total number of species from all sites is used to approximate the total diversity of aquatic plants and the percent of invasive-exotic plants on the lake and in the watershed (Table 2). Many of these plants are considered ecologically harmful, as they tend to out-compete native species. Such “nuisance” plants can also make boating and other recreational activities difficult or impossible. The common and scientific names of plant species found on your lake are listed in Table 3.

Table 2. Comparison of species diversity between your lake and other assessed lakes located within your watershed.

	<u>Lake Keene</u>	<u>Cypress Creek</u> (Average)
Number of Taxa:	28	26
Percent Exotic Plants:	18%	15%

Table 3. Botanical and common names of the most commonly found plants on your lake. Percent frequency (of occurrence), habit (location where found), status (native or exotic), and EPPC status are provided.

<u>Common Name</u>	<u>Plant Species</u>	<u>Frequency</u>	<u>Habit</u>	<u>Status</u>	<u>EPPC</u>
Water Hyacinth	Eichhornia crassipes	100%	Floating	Exotic	I
Water Primroses, Primrosewillow	Ludwigia spp.	100%	Emergent	Unknown	NL
Para Grass	Urochloa (Brachiaria) mutica	90%	Emergent	Exotic	I
Burhead Sedge, Cuban Scirpus	Scirpus cubensis	80%	Emergent	Native	NL
Alligator Weed	Alternanthera philoxeroides	70%	Emergent	Exotic	II
Torpedo Grass	Panicum repens	70%	Emergent	Exotic	I
Water Spangles, Water Fern	Salvinia minima	70%	Floating	Native	NL
Giant Duckweed	Spirodela polyrhiza	70%	Floating	Native	NL
Cypress	Taxodium spp.	70%	Emergent	Native	NL
Manyflower Marshpennywort, Water Penny	Hydrocotyl umbellata	50%	Emergent	Native	NL
Cattails	Typha spp.	50%	Emergent	Native	NL
Southern Red Maple	Acer rubrum var. trilobum	40%	Emergent	Native	NL
Wax Myrtle	Myrica cerifera	40%	Emergent	Native	NL
Baldwin's Spikerush, Roadgrass	Eleocharis baldwinii	30%	Submersed	Native	NL
Algae	Periphyton spp.	30%	Submersed	Native	NL
Spatterdock, Yellow Pondlily	Nuphar lutea var. advena	20%	Floating	Native	NL

# Lake Keene    10/14/98    Watershed: Cypress Creek

Laurel Oak; Diamond Oak	<i>Quercus laurifolia</i>	20%	Emergent	Native	NL
Camphor-tree	<i>Cinnamomum camphora</i>	10%	Emergent	Native	I
Dahoon Holly	<i>Ilex cassine</i>	10%	Emergent	Native	NL
Sweetbay Magnolia	<i>Magnolia virginiana</i>	10%	Emergent	Native	NL
Manatee Mudflower, Baby's Tears	<i>Micranthemum glomeratum</i>	10%	Submersed	Native	NL
Climbing Hempvine	<i>Mikania scandens</i>	10%	Emergent	Native	NL
Cinnamon Fern	<i>Osmunda cinnamomea</i>	10%	Emergent	Native	NL
Pine Tree	<i>Pinus spp.</i>	10%	Emergent	Native	NL
Smartweed, Knotweed	<i>Polygonum spp.</i>	10%	Emergent	Native	NL
Pickrel Weed	<i>Pontederia cordata</i>	10%	Emergent	Native	NL
Willow	<i>Salix spp.</i>	10%	Emergent	Native	NL
Creeping Oxeye	<i>Sphagneticola (Wedelia) trilobata</i>	10%	Emergent	Exotic	II

Standing Crop

In addition to an overall survey of the types of plants on a lake, an estimate of the standing crop (biomass) of the lake has been obtained for many lakes. This was done by calculating the average weight of the vegetation within a quarter-meter square quadrat tossed haphazardly into three zones (see Figure) at each sampling site around the lake: (1) the emergent zone, (2) the floating zone and (3) the submersed zone. The average weight of the plants (Table 4) from all sampling sites and the dominant type of vegetation (Table 5) are provided. If data tables are not shown, no standing crop estimates were obtained for this lake.




Table 4. Comparison between the average biomass from three zones within your lake and among all lakes assessed within your watershed.

	<u>Lake Keene</u>	<u>Cypress Creek</u> (Average)
Emergent Zone:	8.54	4.39
Floating Zone:	6.36	1.24
Submersed Zone:	0.00	0.34

Number of lakes sampled in your watershed: 10

Note: All biomass measurements are shown in kilograms per square meter.

Table 5. Dominant taxa from three zones within your lake.

<u>Zone</u>	<u>Dominant Plant</u>	<u>Status</u>
Emergent Zone:	Torpedo Grass	Exotic
Floating Zone:	Water Hyacinth	Exotic
Submersed Zone:		